

■ CNE Article

Pilates for Breast Cancer Survivors: Impact on Physical Parameters and Quality of Life After Mastectomy

Daniela L. Stan, MD, Sarah M. Rausch, PhD, Kathleen Sundt, RN, PMA[®]-CPT, RYT-200, Andrea L. Cheville, MD, James W. Youdas, PT, David A. Krause, PT, Judy C. Boughey, MD, Molly F. Walsh, DO, Stephen S. Cha, MS, and Sandhya Pruthi, MD



© iStockphoto.com/Deborah Cheramie

Pilates has been advocated for rehabilitation of breast cancer survivors despite little scientific evidence. The authors of this article have examined the feasibility of a Pilates program in postmastectomy breast cancer survivors and the impact on physical and psychological parameters. Fifteen breast cancer survivors were recruited in a one-arm study of 12 weeks of Pilates exercises. The authors assessed recruitment, adherence, and attrition, and measured changes in shoulder and neck range of motion, posture, height, arm volume, quality of life, mood, and body image from pre- to postintervention. Of 26 eligible patients, 15 enrolled, 13 completed the study, and 10 performed more than 50% of the recommended sessions. Statistically significant improvements emerged for shoulder abduction and internal rotation on the affected side, neck rotation toward the unaffected side, and neck flexion. The affected side arm volume and the interlimb volume discrepancy increased. Significant improvements were reported in quality of life, mood, and body image. The improvements in physical and psychological outcomes are promising and deserve further evaluation in a randomized, controlled study. The increase in affected arm volume also warrants additional investigation.

Daniela L. Stan, MD, is a physician in the Breast Diagnostic Clinic in the Division of General Internal Medicine at the Mayo Clinic in Rochester, MN; Sarah M. Rausch, PhD, is the director of Integrative Medicine, Health Outcomes, and Behavior at Moffitt Cancer Center and Research Institute in Tampa, FL; Kathleen Sundt, RN, PMA[®]-CPT, RYT-200, is a Pilates instructor at the Gentle Pilates and Yoga Studio in Boston, MA; Andrea L. Cheville, MD, is an associate professor in Physical Medicine and Rehabilitation, James W. Youdas, PT, is an associate professor in the College of Medicine, David A. Krause, PT, is an assistant professor in Physical Medicine and Rehabilitation, and Judy C. Boughey, MD, is an associate professor in the Department of Surgery, all at the Mayo Clinic in Rochester, MN; Molly F. Walsh, DO, was an instructor in plastic surgery at the Mayo Clinic in Rochester, MN, at the time of this research, and is now a plastic surgeon with the Arizona Plastic and Reconstructive Surgeons in Scottsdale; and Stephen S. Cha, MS, is a statistician III in Health Science Research and Sandhya Pruthi, MD, is a consultant in the Breast Diagnostic Clinic in the Division of General Internal Medicine, both at the Mayo Clinic in Rochester, MN. The authors take full responsibility for the content of the article. This research was funded by grants from Join the Journey and the Ralph and Radene Stump Foundation. The content of this article has been reviewed by independent peer reviewers to ensure that it is balanced, objective, and free from commercial bias. No financial relationships relevant to the content of this article have been disclosed by the independent peer reviewers or editorial staff. Stan can be reached at stan.daniela@mayo.edu, with copy to editor at CJONEditor@ons.org. (First submission May 2011. Revision submitted June 2011. Accepted for publication July 12, 2011.)

Digital Object Identifier:10.1188/12.CJON.131-141

For women in the United States, breast cancer is the leading cancer diagnosis and second most frequent cause of cancer-related deaths. Despite increasing survival rates, the disease and its treatment continue to burden survivors with adverse sequelae such as upper extremity impairment, lymphedema, fatigue, depression, weight gain, and immune system dysfunction (Cheville & Tchou, 2007; Demark-Wahnefried et al., 2001; Karki, Simonen, Malkia, & Selfe, 2005; Rietman et al., 2003). Some studies have indicated that patients who undergo mastectomy experience a lower quality of life (QOL) and more shoulder morbidity compared to patients who undergo lumpectomy (Engel, Kerr, Schlesinger-Raab, Sauer, & Holzel, 2004; Nesvold, Dahl, Lok-

kevik, Mengshoel, & Fossa, 2008; Skrzypulec, Tobor, Drosdzol, & Nowosielski, 2009).

Evidence shows that physical activity in general improves QOL, mood, fatigue, body image, and fitness (Courneya et al., 2007; Galvao & Newton, 2005; Knols, Aaronson, Uebelhart, Franssen, & Aufdemkampe, 2005; McCausland, 2010; McNeely et al., 2006; Ohira, Schmitz, Ahmed, & Yee, 2006; Pinto et al., 2008; Stevinson, Lawlor, & Fox, 2004) and may contribute to primary and secondary breast cancer prevention (Friedenreich, Gregory, Kopciuk, Mackey, & Courneya, 2009; Holmes, Chen, Feskanich, Kroenke, & Colditz, 2005; Peters et al., 2009). Despite this, a structured approach to physical exercise after breast cancer treatment is neither routinely recommended by