

Effect of Isometric Hand Grip Exercises on Blood Flow and Placement of IV Catheters for Administration of Chemotherapy

Ayşe Ozkaraman, PhD, and Öznur Usta Yesilbalkan, PhD



© simm49/iStock/Thinkstock

Background: Complications may occur in the subcutaneous or subdermal tissues during IV administration of chemotherapy related to blood flow and catheter placement.

Objectives: Daily isometric hand grip exercises were evaluated for their effect on blood flow in the vessels of the nondominant arm before placement of IV catheters and the success rate of IV catheter placement on the first attempt.

Methods: The study focused on patients with non-Hodgkin lymphoma receiving the first and second cycles of chemotherapy. The intervention group performed daily isometric hand grip exercises before chemotherapy with peripheral catheter insertion. The control group performed routine activities only. Blood flow was measured by ultrasound in the brachial artery (BA) and brachial vein (BV) of the nondominant arm before the first (T1) and second (T2) cycles of chemotherapy.

Findings: Blood flow slightly increased in the intervention group at T2 compared to T1. In the control group, blood flow decreased in the BA and did not change in the BV at T2 compared to T1. The success rate for first-attempt placement of a peripheral IV catheter was the same for the intervention and control groups.

Ayşe Ozkaraman, PhD, is an assistant professor in the Faculty of Health Sciences at Eskisehir Osmangazi University and Öznur Usta Yesilbalkan, PhD, is an associate professor in the Faculty of Nursing at Ege University in Izmir, both in Turkey. The authors take full responsibility for the content of the article. The authors did not receive honoraria for this work. The content of this article has been reviewed by independent peer reviewers to ensure that it is balanced, objective, and free from commercial bias. No financial relationships relevant to the content of this article have been disclosed by the authors, planners, independent peer reviewers, or editorial staff. Ozkaraman can be reached at aoyaydin26@hotmail.com, with copy to editor at CJONEditor@ons.org. (Submitted March 2015. Revision submitted November 2015. Accepted for publication November 9, 2015.)

Key words: isometric hand grip exercises; blood flow; intravenous chemotherapy

Digital Object Identifier: 10.1188/16.CJON.AE-01

The incidence of hematologic cancers, such as multiple myeloma and leukemia, are increasing, and patients with these cancers are usually treated with systemic chemotherapy (Turkish Society of Hematology Publications, 2011). Although nurses carefully administer chemotherapy, the prevention of all complications, including the extravasation of chemotherapy into the peripheral tissues, is not always possible (Fidalgo et al., 2012; Schulmeister, 2010; WOSCAN Cancer Nursing and Pharmacy Group, 2009). Peripheral IV catheter (PIVC) application, duration, and type of treatment are the major factors that are effective in the success of peripheral IV chemotherapy administration (Centre for Healthcare Related Infection Surveillance and Prevention and Tuberculosis Control, 2013; Fidalgo et al., 2012; Infusion Nurses Society, 2011); however, blood flow at the administration site also is of great importance (Dural, 2008; Türker and Kayaalp, 2002). IV administered chemotherapy drugs rapidly

enter the circulation, followed by dilution on distribution to the systemic plasma (Dural, 2008; Süzer, 2008; Türker & Kayaalp, 2002). Complication risks, such as extravasation in the tissues, increase if dilution of the drug is delayed, particularly in the smaller veins of the arms with slower blood flow (Dural, 2008). In the current study, the effects of isometric hand grip exercises were evaluated to test the hypothesis that an increase in the blood flow velocity would result in faster successful insertion of PIVCs and fewer complications, such as extravasation in the peripheral tissues.

To prevent complications and maintain effective drug delivery, venodilation (e.g., tourniquet application, regional hot application) and veno-constriction (e.g., hand grip exercises) methods have been used to increase the blood flow in the peripheral veins (Ay, 2007; Beer, 2002; Fink et al., 2009; Infusion Nurses Society, 2011; Khawaja, Williams, & Weaver, 1991; Lenhardt, Seybold, Kimberger, Stoiser, & Sessler, 2002;